

CLENPIQ PREP (procedure 10 AM and later)

General Guidelines:

- If CLENPIQ is not covered by your insurance, please use a coupon at <https://www.clenpiq.com/hcp/coverage>
- If you have a **defibrillator**, are on **dialysis**, or are **significantly overweight (BMI>50)**, you must schedule your procedure at a hospital (not an outpatient endoscopy center).
- If you take certain diabetes or weight loss medicines including Ozempic, Rybelsus, Wegovy, Mounjaro, Adlyxin, Trulicity, Victoza, Saxenda, Byetta, or Bydureon, you must stop your medicine at **least 1 week prior** to your colonoscopy.
- For 5 days prior to your colonoscopy, avoid fiber supplements, seeds, nuts, skins of fruits, vegetables, whole grain bars and bread.
- Iron, vitamin E, and NSAIDs (e.g. ibuprofen, naproxen, and diclofenac) should be stopped 3 days prior to your colonoscopy. Tylenol, aspirin, and multivitamins are permitted.
- If you take blood thinners or anti-platelet agents such as Coumadin, Eliquis, Xarelto, Pradaxa, Plavix, Effient or Brilinta, you must contact your **cardiologist/internist** to find out when to stop your medication.
- No marijuana 12 hours prior to the procedure.
- Due to the sedation for your procedure, you **must** be accompanied by a driver. No driving is permitted until the following day.

FOLLOW THESE INSTRUCTIONS (please disregard the manufacturer's instructions):

The day before the colonoscopy—all day:

Follow a **clear liquid diet**. Do not eat any solid food (including breakfast). Please see examples on the back of this sheet. Drink as much fluid as you can throughout the day.

The day before the colonoscopy, at **7 PM:**

1. Take 2 extra strength Gas-X (simethicone) 125mg soft gels or tablets).
2. Drink the entire first bottle of the prep.
3. Drink five (5) 8-ounce drinks of clear liquids (total of 40 ounces).
4. If you become uncomfortable, consume the solution/water at a slower pace and keep walking around to promote intestinal motility.

The day of your colonoscopy (procedure 10 AM and later):

Starting **5 hours** prior to your scheduled procedure, take the second preparation dose:

1. Take 2 extra strength Gas-X (simethicone) 125mg soft gels or tablets).
2. Drink the entire second bottle of the prep.
3. Drink five (5) 8-ounce drinks of clear liquids (total of 40 ounces).
4. If you become uncomfortable, consume the solution/water at a slower pace and keep walking around to promote intestinal motility.
5. Optional: If you do not feel clear after completing the prescribed prep, take 3 capfuls of Miralax in 8 oz of water. You may repeat this until the stool is clear.

You must **complete the entire preparation at least 3 hours** prior to your procedure start time.

Do not drink any liquids (including Miralax) within 3 hours of your procedure or it may be cancelled.

Name: _____ Procedure Date: _____

REGISTER AT: _____ AM. PM. PROCEDURE AT: _____ AM. PM.

REPORT TO:

____ Florham Park Endoscopy Center, 195 Columbia Turnpike, Florham Park, NJ. 973-410-1800

____ Hanover Endoscopy Center of Northern NJ. 91 South Jefferson Rd, Suite 300 Whippany, NJ. 973-929-6800.

____ Cooperman Barnabas Medical Center, Cooperman Family Pavilion, Main Entrance, 94 Old Short Hills Rd, Livingston, NJ 07039. 973-322-5062