

PLENVU PREP (procedure 10 AM and later)**General Guidelines:**

- If PLENVU is not covered by your insurance, please use a coupon at <https://www.myplenvu.com/resources/>
- If you have a **defibrillator**, are on **dialysis**, or are **significantly overweight (BMI>50)**, you must schedule your procedure at a hospital (not an outpatient endoscopy center).
- If you take certain diabetes or weight loss medicines including Ozempic, Rybelsus, Wegovy, Mounjaro, Adlyxin, Trulicity, Victoza, Saxenda, Byetta, or Bydureon, you must stop your medicine at least **8 days prior** to your colonoscopy.
- For 5 days prior to your colonoscopy, avoid fiber supplements, seeds, nuts, skins of fruits and vegetables, whole grain bread and bars.
- Iron, vitamin E, and NSAIDs (e.g. ibuprofen, naproxen, and diclofenac) should be stopped 3 days prior to your colonoscopy. Tylenol, aspirin, and multivitamins are permitted.
- If you take blood thinners or anti-platelet agents such as Coumadin, Eliquis, Xarelto, Pradaxa, Plavix, Effient or Brilinta, you must contact your **cardiologist/internist** to find out when to stop your medication.
- No marijuana 12 hours prior to the procedure.
- Due to the sedation for your procedure, you **must** be accompanied by a driver. No driving is permitted until the following day.

FOLLOW THESE INSTRUCTIONS (please disregard the manufacturer's instructions):**The day before the colonoscopy—all day:**

Follow a **clear liquid diet**. Do not eat any solid food (including breakfast). Drink as much fluid as you can throughout the day.

The day before the colonoscopy, at 7 PM:

1. Take 2 extra strength Gas-X (simethicone) 125mg soft gels or tablets).
2. Open the "Dose 1" packet and pour the powder into the provided mixing container. Fill the container with 16oz of water and mix until completely dissolved. Drink the entire solution within 30 minutes.
3. Fill the container with another 16oz of water and drink within 60 minutes. If you become uncomfortable, consume the solution/water at a slower pace and keep walking around to promote intestinal motility.
4. You may keep drinking clear liquids until midnight.

The day of your colonoscopy (procedure 10AM and later):

Starting **5 hours** prior to your scheduled procedure, take the second preparation dose:

1. Take 2 extra strength Gas-X (simethicone) 125mg soft gels or tablets).
2. Open the "Dose 2" (A and B) packets and pour the powders into the provided mixing container. Fill the container with 16oz of water and mix until completely dissolved. Drink the entire solution within 30 minutes.
3. Fill the container with another 16oz of water and drink within 60 minutes.
4. Optional: If you do not feel clear after completing the prescribed prep, take 3 capfuls of Miralax in 8 oz of water. You may repeat this until the stool is clear.

You must **complete the entire preparation at least 3 hours** prior to your procedure start time.
Do not drink any liquids (including Miralax) within 3 hours of your procedure or it may be canceled.

Name: _____ Procedure Date: _____

REGISTER AT: _____ AM. PM. PROCEDURE AT: _____ AM. PM.

REPORT TO:

____ Florham Park Endoscopy Center, 195 Columbia Turnpike, Florham Park, NJ. 973-410-1800

____ Hanover Endoscopy Center of Northern NJ. 91 South Jefferson Rd, Suite 300 Whippany, NJ. 973-929-6800.

____ Cooperman Barnabas Medical Center, Cooperman Family Pavilion, Main Entrance, 94 Old Short Hills Rd, Livingston, NJ 07039. 973-322-5062

____ Morristown Surgical Center, 111 Madison Ave, 4th floor, Morristown, NJ 07960, 973-971-6970

____ Morristown Medical Center, 100 Madison Ave, Morristown, NJ 07960, Level D, Digestive Health Department/
Goryeb Children Hospital entrance 973-971-5088